

3° 3âµ j µ*(' © šµ- µ®qj'
fi m©l . » ' ' « - , - fl

3° 3âµ ç µ*(' ©šµ- µ®qj' w-m©. » ' ' « - , - » j ' 2° 3° 3âµ j µ° 3° çñ*(' ©j' ©â ' mō³ ° » 3 TM° @k a l »'
3 » ' - - - ° » 2 ' ' TM° - - ± . , µ ' ©' - - ° ' - 3 ' ' - - ©' » j ' n a l - - ° ' 2 »ñ ® l »' Q

q@3 TM°

- ° µ'	- - - - - 2	©' » â µ³ ' ' 2		® - - - - 2 fjµ α' fl	m ô©² ' Ÿ j - - - ©ª ' - - ° µ' - - - - ° œ mō©' µ' fi« - fl
		° !' ©ñ 2	° ' œ- ' 2 " ©' 2 !µ !		
- -	° » ° 2 ' °	225-300	300-400	500	30

q@3 TM° - µ w - « , ' °

® - ° ' ¼ 3 ' - ž
® ¥ ®' µ « - ' µ' Ÿ j - - ° 3 ' j ° ®' µ - - - - j µ' µ™ - - ° 3 TM° - 3 ' j w Q

3 TM° © l ' - µ w ° . n - 3 ' °

- 1! ' - - ° 2 ' ž - - ° 2 ' - - - µ ' ° © l | ' ' ®, { ' - - œ ' ° ' « ' ' ' Q
- 2! ' 2 » ž ñ œ | ' ' n - - - ° ú @ - l ° ' ° œ³ Q
- 3! ' Ÿ j - - - - ° ® - - ° ° ' ' mōk ' j - - - ° œ³ Q » ' - - « , ' 2 ' Ÿ j - - - - Q
- 4! ' Ÿ j - - - - ° « ' « 1 ©' - ® j | ' ' ' - - m™ - - mç Ÿ ' ©' » - w Q
- 5! ' Ÿ j - - - - ° 2 3 ' - ® - ž - - ž ® - | ' - Ÿ ' œ' - - - » ' œ » w Q
- 6! ' Ÿ j - - - - ° 3 ' 2 µ © ' ° 2 3 ' ® l ° ' ¼ ñ° - - ® l ' ® » - Q

. 1 ' - µ ¼ ' °

° ' « ž œâ - - n - ž² ©µ ž qý µ ž ç - - f j² ' TM fl » ' ° - © ' » Q

a 2 - - œ - ñ°

- 1! ' 3 « ' - TM µ j 3 ' © ' TM µ - - ® Ÿ ' TM k TM k - - - qý µ - - 3 ' 3 » ' 3 ' © ' © - « » ' © ' j ' ° © - qý µ - - - µ ' ® k a l ° - - » ' j w Q 3 « ' 2 ' j ' ° » , ' » ' © ' qý µ - - - 3 Q
- 2! ' 3 « ' - ® j | ' ' ñ œ « 1 ©' » j w © - - Ë' q © ' « | ' ' « 1 ©' ñ œ - - - - 2 ' 2 ' ° ° - | ' ® - - ° ' - w Q
- 3! ' 3 « ' n - - « 1 ©' » j 3 ' © ' q - - - - 2 ' 2 ' ° µ o - - ° ® - - ° µ TM ± TM 10-15 ' 2 - - - - w Q
- 4! ' 3 « ' ° ° ' ' mōk ' TM B ' » ' © ' TM - - , ' » . ' 2 ' µ j ' 3 Q

® - , - ©

° ç . - - 2 ' - m - ° ' 3 TM - - - ® ' 3 » ® - , - - - » ' » Q

. 1 ' - - -

Caution: "Not to Be Used on Crops Other Than Specified on This Label/Leaflet".

Tricyclazole 75% W.P.**(Systemic Fungicide)**

Tricyclazole 75% WP is a systemic fungicide based on the active ingredient, Tricyclazole 75% w/w and balance adjuvants.

Recommendation

Crop(s)	Common Name of Pest	Dosage/HA		Dilution in Water (Liter)	Waiting Period between last spray to harvest (Days)
		AI gm	Formulation gm/ml.		
Paddy	Blast	225-300	300-400	500	30

Direction of Use

Plant Protection equipment :
knapsack sprayer fitted with hollow cone nozzle

Precaution

1. Keep away from foodstuffs, empty foodstuff containers and animals food.
2. Avoid contact with mouth, eyes and skin.
3. Avoid inhalation the spray mist. Spray in the direction of wind.
4. Wash thoroughly the contaminated clothes and parts of the body after spraying.
5. Do not smoke, drink, eat and chew anything while spraying.
6. Wear full protective clothing while mixing and spraying.

Symptoms Of Poisoning

Headache, giddiness, nausea, vomiting, convulsions may occur.

First Aid

1. If swallowed, induce vomiting by tickling the back of throat. Repeat it until the vomitus is clear. Do not induce vomiting if the patient is unconscious.
2. If clothing and skin are contaminated, remove the clothes and wash the contaminated skin with copious amount of soap and water.
3. If eyes are contaminated, flush with plenty of saline/clean water for about 10 to 15 minutes.
4. If inhaled, remove the patient to fresh air.

Phytotoxicity

The product is not phytotoxic when used as per field recommendation.

Antidote

No specific antidote is known. Treat symptomatically.

Disposal Of Used Container

- 1 It shall be the duty of manufactures formulators and operator to dispose packages or surplus materials and washings from the machine and container shall be disposed off in a safe manner so as to prevent environmental and water pollution.
- 2 The used packages shall not be left outside to prevent their re-use.
- 3 Packages shall be broken and buried away from habitation.

Storage Conditions

Caution: "Not to Be Used on Crops Other Than Specified on This Label/Leaflet".

1. The packages containing the fungicides shall be stored in separate rooms or premises away from the rooms or premises used for storing other articles particularly articles of or shall be kept in separate almirahs under lock and key depend upon the quantity and nature of the fungicide.

2. The store rooms should be well built, dry, well-lit and ventilated and of sufficient dimensions. The contaminated of the store should be dry and cool.

Chemical Composition:

Tricyclazole technical	(min purity 95% w/w a.i.)	78.90 % w/w
Sodium Lignosulphonate		3.00 % w/w
Sodium Lauryl Sulphonate		3.00 % w/w
Hisil	(Synthetic hydrated precipitated silica)	5.00 % w/w
China Clay		10.10 % w/w
		Total: 100.000% w/w

Manufactured By:

Toshi Insecticides India
Gaushala Road, Near Cooler Factory, Geeta House, Karnal

Karnal, Karnal, Haryana

Manufacturer Premises :

Meerut Road, Village Andhera, District-Karnal, Haryana-132001

Caution: "Not to Be Used on Crops Other Than Specified on This Label/Leaflet".