

—'x®» o âµ' o' (#' ©' ©' w' !®!

fi'' —x— , —fl

3 » w—šµ—, µ'®q!''—x—, —»' j ° 2 ± '—° '(#' ©' ©' 3'©ñ' —'x®» o âµ' o'!®'° —'mô'° »3 TM'®« a | »'Q—'x®
 » o âµ' o'!®' (#' ©'! w' !®! w—° »©' m° «' —'x—, —»' j ° —'q®3 TM' —' —'©—'Y«—'®' !· µ— —'x— —' —'a 2' »©' —'3 '
 j' ©' »'Q o° —'q®3 TM' —'©' —'«'° —'x—, —'° ±ô—' »'Q 3'—'©—'· µ'®'æ' —'· µ'—x—' µw m° «' ' » | —'x—' —'ú—
 ° '©' 2' »' —'3 —' —'©' »'Q 3'—'x—, —'x—' —'° ±' m— a { ' 2' —'3 —' —'©' »'®'° —' —' ±' «' —' µ'—' »'Q —'x—' —'©—
 « 2 —'®'° —'m° —' —'» »'©' »'Q | —'©—' 3' —' µw ŷ³ · »' —'®'° · 1 µ'—' »' »'Q

q®3 TM'

° µ'	—x'—' 2	©'»â x³' 2		®—'—' 2 fµ x'fl	m ô®²' Y'—', ®'° —'° µ'—x—'— ° æ' mô®' µ' fi«—fl
		°'©ñ fi ² fl	°'æ—' fi—! !fl		
æ· µ	©—' Y«— ®' !· µ—	500	1	500-1000	21
		500	1	500-1000	21

q®3 TM' —' µ w— «'—

q®3 TM' 2' n —'· µ' 3' —'®—' 3' Z' —'x' 3' Z' —'ú—, —'—'®—' —'° x'—' 3' ' | —' w' !®!p!p!žvôd®!®!·®'° 3' ' n «' Q

3 TM'© { —' µ w° · n —'3' }

- 1! —' 2 Z' —' 2 —' —' µ' —'©'—' | —'®, { —'æ' —'° «' —'° Q
- 2! 2 » Z' ñ æ' | —' n —' —'° ú®—' —'° æ³ Q
- 3! Y Ê—' —' —' ®' —'° · —' —' mō«' j' —' —'° æ³ Q» —' —' «' 2' Y Ê—' —' —' Q
- 4! Y Ê—' —' —' «' «' 1©—'® | —' , —' —' m™ —' mçY '©' » n wQ
- 5! Y Ê—' —' —' 2 3' n ®—' Z' —' Z' ®—' | —' —'Y 'æ' —' —' »æ » wQ
- 6! Y Ê—' —' —' 3' 2 µ' ©' 2 3' ®' | ° ¼ ñ² —' —'® | ®» —' Q

· 1' —' µ ¼'

2 ©µ Z' , —' w' —'° · 2 —'®—' Z' µ' Z' ® | 2 Z' | ® —'3 | —' 2 o 3 ° · »' —'° —'® » Q

a 2 —' æ—'ñ'

- 1! 3 «' —'™µ' j' 3'©'™µ' —'®'Y'™µ'™µ' —' —' qy x' —' 3' 3' »' 3'©'° —' «' »'©' »' j' —'©' —' qy x' · —' —' µ' —'®«' a | ° —' —' »' j' wQ
- 3 «' 2' j' —' »' »'©' qy x' —' —' 3' Q
- 2! 3 «' —'® | —' ñ æ' «' 1©' »' j' w' ©' «' 1©' —'®—' q®' «' —' | —' ñ æ' —' —' 2' 2'° · —' | —'®—' —' n wQ
- 3! 3 «' n —' «' 1©' »' j' 3'©' q—' —' —' 2' 2'° µ' o—'° —' —'®' —'° µ'™±™™ 10-15' 2' —'x' —' —' wQ
- 4! 3 «' ° · —' —' mō«' —'™ 3' »'©' —'™ —' —' , —' »' 2' µ' j' 3' Q

®—' · , â©©

Caution: Í Not to Be Used on Crops Other Than Specified on This Label/Leaflet! "

Cartap Hydrochloride 50% SP**(Insecticide)**

It is a soluble insecticide which contains 50% Cartap Hydrochloride active ingredient and balance adjuvants. Cartap hydrochloride 50% S.P. is an insecticide highly effective for the control of stem borer and leaf folder on rice. Cartap Hydrochloride has a quite different mode of action from that of other insecticides. It is effective against both chewing and sucking insects and controls at a low concentration. Cartap hydrochloride controls almost all stages of insects and the efficacy of Cartap hydrochloride is prolonged. It has no effect to natural enemies of insect pests and it is practically nontoxic to mammals

Recommendation

Crop(s)	Common Name of Pest	Dosage/HA		Dilution in Water in water (liter)	Waiting Period between last spray to harvest days
		AI (gm)	Formulation (kg)		
Rice	Stem borer	500	1	500-1000	21
	Leaf Folder	500	1	500-1000	21

Direction of Use

P.P. Equipment:- Knapsack sprayer, foot sprayer,compression knapsack battery sprayer and ASPEE-HTP power sprayer.

Precaution

1. Keep away from foodstuffs, empty foodstuff containers and animals food.
2. Avoid contact with mouth, eyes and skin.
3. Avoid inhalation the spray mist. Spray in the direction of wind.
4. Wash thoroughly the contaminated clothes and parts of the body after spraying.
5. Do not smoke, drink, eat and chew anything while spraying.
6. Wear full protective clothing while mixing and spraying.

Symptoms Of Poisoning

Nausea, trembling of extremities of body, salivation, spasms, dyspnoea and mydriasis may occur.

First Aid

1. If swallowed, induce vomiting by tickling the back of throat. Repeat it until the vomitus is clear. Do not induce vomiting if the patient is unconscious.
2. If clothing and skin are contaminated, remove the clothes and wash the contaminated skin with copious amount of soap and water.
3. If eyes are contaminated, flush with plenty of saline/clean water for about 10 to 15 minutes.
4. If inhaled, remove the patient to fresh air.

Phytotoxicity

Cartap hydrochloride 50% SP is not phytotoxic to crop at recommended dosage

Antidote

An intravenous injection of 100-200 mg of L-cysteine or an intramuscular injection of 20-60 mg of BAL (dimercaprol or 2,3 dimercapatoproponal) are recommended. Apply steroids against dermatitis.

Disposal Of Used Container

Caution: Not to Be Used on Crops Other Than Specified on This Label/Leaflet

1. Package surplus materials and washing from the machines and containers should be disposed or in a safe manners of sufficient prevent environmental or water pollution.
2. The used package shall not be left outside to prevent their re-use
3. Package shall be broken and burried away from habitation

Storage Conditions

1. The packages containing the insecticides shall be stored in separate room or premises away from the rooms or premises used for storing other articles particularly food or shall be kept in separate almirahs under lock and key depending upon the quantity and nature of the insecticides
2. The room or premises menat for storing the insecticides shall be well built, dry well lit, and ventilated and of sufficient dimension to avoid contamination with vapour

Chemical Composition:

Cartap Hydro chloride Tech.	(Based on 98% w/w)	51.20 % w/w
Surfactant	(Alkyl phenol ethylene oxide)	0.70 % w/w
Stabilizer	(Isopropyl acid phosphate)	0.30 % w/w
Acid green dye		0.05 % w/w
Diluent	(4-O-B-D gatacto pyronosyl alpha -D glucophyranose/lactose)	47.75 % w/w
Total:		100.000% w/w

Manufactured By:

Toshi Insecticides India
Gaushala Road, Near Cooler Factory, Geeta House, Karnal

Karnal, Karnal, Haryana

Manufacturer Premises :

Meerut Road, Village Andhera, District-Karnal, Haryana-132001

Caution:Í Not to Be Used on Crops Other Than Specified on This Label/Leafletî ¨